

**Patient Instruction Sheet
BUSPIRONE HYDROCHLORIDE, USP**


Rx Only

**HOW TO USE:
Buspirone Hydrochloride Tablets, USP
15 mg and 30 mg Tablets**
in convenient adjustable dosage tablet form


Response to buspirone varies among individuals. Your physician may find it necessary to adjust your dosage to obtain the proper response.

This adjustable dosage tablet design makes dosage adjustments easy. Each tablet is scored and can be broken accurately to provide any of the following dosages.


If your doctor
prescribed
the 30-mg
tablet:




30 mg
(the entire
tablet)



20 mg
(two thirds
of a tablet)



10 mg
(one third
of a tablet)




15 mg
(one half
of a tablet)


If your doctor
prescribed
the 15-mg
tablet:




15 mg
(the entire
tablet)



10 mg
(two thirds
of a tablet)




5 mg
(one third
of a tablet)



7.5 mg
(one half
of a tablet)

To break a adjustable dosage tablet accurately and easily, hold the tablet between your thumbs and index fingers close to the appropriate tablet score (groove) as shown in the photo. Then, with the tablet score facing you, apply pressure and snap the tablet segments apart (segments breaking incorrectly should not be used).



Manufactured by: Oxford Pharmaceuticals LLC
301 Leaf Lake Parkway, Birmingham, AL 35211 USA

8200016
06/20
R01